

wherein said composition comprises less than 4% of an anionic, zwitterionic or amphoteric surfactant and from about 0.01% to about 1.0% of C₁₀ to C₁₆ mono, -di and -tri acyl sugar

elsim 3. (amended)

A composition according to Claim 2 wherein the polyhydric alcohol is selected from the group consisting of glycerine, butylene glycol, propylene glycol, dipropylene glycol, polyethylcne glycol, hexane tripl, ethoxylated glycerine, propoxylated glycerine and mixtures thereof.

Claim 4. (amended)

A composition according to Claim 3 wherein the polyhydric alcohol is selected from the group consisting of glycerine; polyethylene glycol and mixtures thereof.

Claim 6. (amended)

A composition according to Claim 5 wherein the cationic polymer is a cationic polysaccharide selected from the group consisting of cationic cellulosic polymers, cationic polysaccharide gums and mixtures thereof.

Claim 11. (amended)

A cosmetic method for moisturizing skin comprising the step of applying to the skin a composition according to Claim 10.

Claim 12, (amended)

The cosmetic method of claim 11 wherein the composition comprises a cation containing polymer selected from the group consisting of cationic, basic, amphoteric and zwitterionic polymers, and mixtures thereof.

Claim 13 (amended)

The cosmetic method of claim 13 wherein said composition is left on the skin.